

INBRED MAIZE LINE NP2052

FIELD OF THE INVENTION

This invention is in the field of maize breeding, specifically relating to an inbred maize line designated NP2052.

BACKGROUND OF THE INVENTION

The goal of plant breeding is to combine in a single variety or hybrid various desirable traits. For field crops, these traits may include resistance to diseases and insects, tolerance to heat and drought, reducing the time to crop maturity, greater yield, and better agronomic quality. With mechanical harvesting of many crops, uniformity of plant characteristics such as germination and stand establishment, growth rate, maturity, and plant and ear height, is important.

Field crops are bred through techniques that take advantage of the plant's method of pollination. A plant is self-pollinated if pollen from one flower is transferred to the same or another flower of the same plant. A plant is cross-pollinated if the pollen comes from a flower on a different plant. Plants that have been self-pollinated and selected for type for many generations become homozygous at almost all gene loci and produce a uniform population of true breeding progeny. A cross between two different homozygous lines produces a uniform population of hybrid plants that may be heterozygous for many gene loci. A cross of two plants each heterozygous at a number of gene loci will produce a population of hybrid plants that differ genetically and will not be uniform.

Maize (*Zea mays* L.), often referred to as corn in the United States, can be bred by both self-pollination and cross-pollination techniques. Maize has separate male and female flowers on the same plant, located on the tassel and the ear, respectively. Natural pollination occurs in maize when wind blows pollen from the tassels to the silks that protrude from the tops of the ears.

A reliable method of controlling male fertility in plants offers the opportunity for improved plant breeding. This is especially true for development of maize hybrids, which relies upon some sort of male sterility system. There are several options for controlling male fertility available to breeders, such as: manual or mechanical emasculation (or detasseling), cytoplasmic male sterility, genetic male sterility, gametocides and the like.

Hybrid maize seed is typically produced by a male sterility system incorporating manual or mechanical detasseling. Alternate strips of two maize inbreds are planted in a field, and the pollen-bearing tassels are removed from one of the inbreds (female). Providing that there is sufficient

isolation from sources of foreign maize pollen, the ears of the detasseled inbred will be fertilized only from the other inbred (male) and the resulting seed is therefore hybrid and will form hybrid plants.

The laborious, and occasionally unreliable, detasseling process can be avoided by using cytoplasmic male-sterile (CMS) inbreds. Plants of a CMS inbred are male sterile as a result of factors resulting from the cytoplasmic, as opposed to the nuclear, genome. Thus, this characteristic is inherited exclusively through the female parent in maize plants, since only the female provides cytoplasm to the fertilized seed. CMS plants are fertilized with pollen from another inbred that is not male-sterile. Pollen from the second inbred may or may not contribute genes that make the hybrid plants male-fertile. Seed from detasseled fertile maize and CMS produced seed of the same hybrid can be blended to insure that adequate pollen loads are available for fertilization when the hybrid plants are grown.

There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, as disclosed in U.S. Pat. Nos. 4,654,465 and 4,727,219 and chromosomal translocations as described in U.S. Pat. Nos. 3,861,709 and 3,710,511, the disclosures of which are specifically incorporated herein by reference. There are many other methods of conferring genetic male sterility in the art, each with its own benefits and drawbacks. These methods use a variety of approaches such as delivering into the plant a gene encoding a cytotoxic substance associated with a male tissue specific promoter or an antisense system in which a gene critical to fertility is identified and an antisense to that gene is inserted in the plant (see: publications EPO 89/3010153.8 and WO 90/08828).

Another system useful in controlling male sterility makes use of gametocides. Gametocides are not a genetic system, but rather a topical application of chemicals. These chemicals affect cells that are critical to male fertility. The application of these chemicals affects fertility in the plants only for the growing season in which the gametocide is applied (see Carlson, Glenn R., U.S. Pat. No. 4,936,904, which is incorporated herein by reference). Application of the gametocide, timing of the application and genotype specificity often limit the usefulness of the approach.

The use of male sterile inbreds is but one factor in the production of maize hybrids. The development of maize hybrids requires, in general, the development of homozygous inbred lines, the crossing of these lines, and the evaluation of the crosses. Pedigree breeding and recurrent selection breeding methods are used to develop inbred lines from breeding populations. Breeding programs combine the genetic backgrounds from two or more inbred lines or various other germplasm sources into breeding pools from which new inbred lines are developed by selfing and selection of desired phenotypes. The new inbreds are crossed with other inbred lines and the hybrids from these crosses

are evaluated to determine which of those have commercial potential. Plant breeding and hybrid development are expensive and time-consuming processes.

Pedigree breeding starts with the crossing of two genotypes, each of which may have one or more desirable characteristics that is lacking in the other or which complements the other. If the two original parents do not provide all the desired characteristics, other sources can be included in the breeding population. In the pedigree method, superior plants are selfed and selected in successive generations. In the succeeding generations the heterozygous condition gives way to homogeneous lines as a result of self-pollination and selection. Typically in the pedigree method of breeding five or more generations of selfing and selection is practiced: F1 to F2; F3 to F4; F4 to F5, etc.

A single cross maize hybrid results from the cross of two inbred lines, each of which has a genotype that complements the genotype of the other. The hybrid progeny of the first generation is designated F1. In the development of commercial hybrids only the F1 hybrid plants are sought. Preferred F1 hybrids are more vigorous than their inbred parents. This hybrid vigor, or heterosis, can be manifested in many polygenic traits, including increased vegetative growth and increased yield.

The development of a maize hybrid involves three steps: (1) the selection of plants from various germplasm pools for initial breeding crosses; (2) the selfing of the selected plants from the breeding crosses for several generations to produce a series of inbred lines, which, although different from each other, breed true and are highly uniform; and (3) crossing the selected inbred lines with different inbred lines to produce the hybrid progeny (F1). During the inbreeding process in maize, the vigor of the lines decreases. Vigor is restored when two different inbred lines are crossed to produce the hybrid progeny (F1). An important consequence of the homozygosity and homogeneity of the inbred lines is that the hybrid between a defined pair of inbreds will always be the same. Once the inbreds that give a superior hybrid have been identified, the hybrid seed can be reproduced indefinitely as long as the homogeneity of the inbred parents is maintained.

A single cross hybrid is produced when two inbred lines are crossed to produce the F1 progeny. A double cross hybrid is produced from four inbred lines crossed in pairs ($A \times B$ and $C \times D$) and then the two F1 hybrids are crossed again $(A \times B) \times (C \times D)$. Much of the hybrid vigor exhibited by F1 hybrids is lost in the next generation (F2). Consequently, seed from hybrids is not used for planting stock.

Hybrid seed production requires elimination or inactivation of pollen produced by the female parent. Incomplete removal or inactivation of the pollen provides the potential for self-pollination. This inadvertently self-pollinated seed may be unintentionally harvested and packaged with hybrid seed. Once the seed is planted, it is possible to identify and select these self-pollinated plants. These self-pollinated plants will be genetically equivalent to the female inbred line used to produce the

hybrid. Typically these self-pollinated plants can be identified and selected due to their decreased vigor. Female selfs are identified by their less vigorous appearance for vegetative and/or reproductive characteristics, including shorter plant height, small ear size, ear and kernel shape, cob color, or other characteristics.

Identification of these self-pollinated lines can also be accomplished through molecular marker analyses. See, "The Identification of Female Selfs in Hybrid Maize: A Comparison Using Electrophoresis and Morphology", Smith, J. S. C. and Wych, R. D., *Seed Science and Technology* 14, pp. 1-8 (1995), the disclosure of which is expressly incorporated herein by reference. Through these technologies, the homozygosity of the self-pollinated line can be verified by analyzing allelic composition at various loci along the genome. Those methods allow for rapid identification of the invention disclosed herein. See also, "Identification of Atypical Plants in Hybrid Maize Seed by Postcontrol and Electrophoresis" Sarca, V. et al., *Probleme de Genetica Teoritca si Aplicata* Vol. 20 (1) p. 29-42.

As is readily apparent to one skilled in the art, the foregoing are only two of the various ways by which the inbred can be obtained by those looking to use the germplasm. Other means are available, and the above examples are illustrative only.

Maize is an important and valuable field crop. Thus, a continuing goal of plant breeders is to develop high-yielding maize hybrids that are agronomically sound based on stable inbred lines. The reasons for this goal are obvious: to maximize the amount of grain produced with the inputs used and minimize susceptibility of the crop to pests and environmental stresses. To accomplish this goal, the maize breeder must select and develop superior inbred parental lines for producing hybrids. This requires identification and selection of genetically unique individuals that occur in a segregating population. The segregating population is the result of a combination of crossover events plus the independent assortment of specific combinations of alleles at many gene loci that results in specific genotypes. The probability of selecting any one individual with a specific genotype from a breeding cross is infinitesimal due to the large number of segregating genes and the unlimited recombinations of these genes, some of which may be closely linked. However, the genetic variation among individual progeny of a breeding cross allows for the identification of rare and valuable new genotypes. These new genotypes are neither predictable nor incremental in value, but rather the result of manifested genetic variation combined with selection methods, environments and the actions of the breeder. Thus, even if the entire genotypes of the parents of the breeding cross were characterized and a desired genotype known, only a few, if any, individuals having the desired genotype may be found in a large segregating F2 population. Typically, however, neither the genotypes of the breeding cross parents nor the desired genotype to be selected is known in any detail. In addition, it is not

known how the desired genotype would react with the environment. This genotype by environment interaction is an important, yet unpredictable, factor in plant breeding. A breeder of ordinary skill in the art cannot predict the genotype, how that genotype will interact with various climatic conditions or the resulting phenotypes of the developing lines, except perhaps in a very broad and general fashion. A breeder of ordinary skill in the art would also be unable to recreate the same line twice from the very same original parents, as the breeder is unable to direct how the genomes combine or how they will interact with the environmental conditions. This unpredictability results in the expenditure of large amounts of research resources in the development of a superior new maize inbred line.

SUMMARY OF THE INVENTION

According to the invention, there is provided a novel inbred maize line, designated NP2052. This invention thus relates to the seeds of inbred maize line NP2052, to the plants of inbred maize line NP2052, and to methods for producing a maize plant by crossing the inbred line NP2052 with itself or another maize line. This invention further relates to hybrid maize seeds and plants produced by crossing the inbred line NP2052 with another maize line.

The invention is also directed to inbred maize line NP2052 into which one or more specific, single gene traits, for example transgenes, have been introgressed from another maize line. Preferably, the resulting line has essentially all of the morphological and physiological characteristics of inbred maize line of NP2052, in addition to the one or more specific, single gene traits introgressed into the inbred, preferably the resulting line has all of the morphological and physiological characteristics of inbred maize line of NP2052, in addition to the one or more specific, single gene traits introgressed into the inbred. The invention also relates to seeds of an inbred maize line NP2052 into which one or more specific, single gene traits have been introgressed and to plants of an inbred maize line NP2052 into which one or more specific, single gene traits have been introgressed. The invention further relates to methods for producing a maize plant by crossing plants of an inbred maize line NP2052 into which one or more specific, single gene traits have been introgressed with themselves or with another maize line. The invention also further relates to hybrid maize seeds and plants produced by crossing plants of an inbred maize line NP2052 into which one or more specific, single gene traits have been introgressed with another maize line.

DEFINITIONS

In the description and examples that follow, a number of terms are used herein. In order to provide a clear and consistent understanding of the specification and claims, including the scope to be given such terms, the following definitions are provided. Below are the descriptors used in the data tables included herein. All linear measurements are in centimeters unless otherwise noted.

Heat units	$(\text{Max Temp}(\leq 86 \text{ deg. F.}) + \text{Min Temp}(\geq 50 \text{ deg. F.})) / 2 - 50$
EMRGN	Final number of plants per plot
KRTP	Kernel type: 1.sweet 2.dent 3.flint 4.flour 5.pop 6.ornamental 7.piecorn 8.other
ERTLP	% Root lodging (before anthesis)
GRNSP	% Brittle snapping (before anthesis)
TBANN	Tassel branch angle of 2nd primary lateral branch (at anthesis)
LSPUR	Leaf sheath pubescence of second leaf above the ear (at anthesis) 1-9 (1=none)
ANGBN	Angle between stalk and 2nd leaf above the ear (at anthesis)
CR2L	Color of 2nd leaf above the ear (at anthesis)
GLCR	Glume Color
GLCB	Glume color bars perpendicular to their veins (glume bands): 1.absent 2.present
ANTC	Anther color
PLQUR	Pollen Shed: 0-9 (0=male sterile)
HU1PN	Heat units to 10% pollen shed
HUPSN	Heat units to 50% pollen shed
SLKC	Silk color
HU5SN	Heat units to 50% silk
SLK5N	Days to 50% silk in adapted zone
HU9PN	Heat units to 90% pollen shed
HUPLN	Heat units from 10% to 90% pollen shed
DA19	Days from 10% to 90% pollen shed
LAERN	Number of leaves above the top ear node
MLWVR	Leaf marginal waves: 1-9 (1=none)
LFLCR	Leaf longitudinal creases: 1-9 (1=none)
ERLLN	Length of ear leaf at the top ear node
ERLWN	Width of ear leaf at the top ear node at the widest point
PLHCN	Plant height to tassel tip
ERHCN	Plant height to the top ear node

LTEIN	Length of the internode between the ear node and the node above
LTASN	Length of the tassel from top leaf collar to tassel tip
LTBRN	Number of lateral tassel branches that originate from the central spike
EARPN	Number of ears per stalk
APBRR	Anthocyanin pigment of brace roots: 1.absent 2.faint 3.moderate 4.dark
TILLN	Number of tillers per plant
HSKC	Husk color 25 days after 50% silk (fresh)
HSKD	Husk color 65 days after 50% silk (dry)
HSKTR	Husk tightness 65 days after 50% silk: 1-9 (1=loose)
HSKCR	Husk extension: 1.short (ear exposed) 2.medium (8 cm) 3.long (8-10 cm) 4.very long (>10 cm)
HEPSR	Position of ear 65 days after 50% silk: 1.upright 2.horizontal 3.pendent
STGRP	% Staygreen at maturity
DPOPN	% dropped ears 65 days after anthesis
LRTRN	% root lodging 65 days after anthesis
HU25	Heat units to 25% grain moisture
HUSG	Heat units from 50% silk to 25% grain moisture in adapted zone
DSGM	Days from 50% silk to 25% grain moisture in adapted zone
SHLNN	Shank length
ERLNN	Ear length
ERDIN	Diameter of the ear at the midpoint
EWGTN	Weight of a husked ear (grams)
KRRWR	Kernel rows: 1.indistinct 2.distinct
KRNAR	Kernel row alignment: 1.straight 2.slightly curved 3.curved
ETAPR	Ear taper: 1.slight 2.average 3.extreme
KRRWN	Number of kernel rows
COBC	Cob color
COBDN	Diameter of the cob at the midpoint
KRTP	Endosperm type: 1.sweet 2.extra sweet 3.normal 4.high amylose 5.waxy 6.high protein 7.high lysine 8.super sweet 9.high oil 10.other
KRCL	Hard endosperm color
ALEC	Aleurone color
ALCP	Aleurone color pattern: 1.homozygous 2.segregating
KRLNN	Kernel length (mm)

KRWDN	Kernel width (mm)
KRDPN	Kernel thickness (mm)
K100N	100 kernel weight (grams)
KRPRN	% round kernels on 13/64 slotted screen
GRLSR	Grey leaf spot severity rating; 1=resistent, 9=susceptible.
INTLR	Intactness rating of plants at time of harvest; 1=all foliage intact, 9=all plants broken below the ear.
LRTLPL	Percentage of plants lodged, leaning >30 degrees from vertical, but unbroken at harvest.
MST_P	Percent grain moisture at harvest.
SCLBR	Southern corn leaf blight severity rating; 1=resistent, 9=susceptible.
STKLP	Percentage of plants with stalks broken below the ear at time of harvest.
YBUAN	Grain yield expressed as bushels per acre adjusted to 15.5% grain moisture.
STBWR	Stewart Bacterial Wilt
ERLNN	Ear Length
CRSTR	Common Rust Rating
GRQUR	Grain Quality
PLTAR	Plant Appearance
HUBLN	Heat Units to Black Layer
TSTWN	Test Weight in LBS/BU
PSTSP	Push Test for Stalk/Root Quality on Erect Plants
ERGRR	Early Growth:6+ Leaf Stage

DETAILED DESCRIPTION OF THE INVENTION

Inbred maize lines are typically developed for use in the production of hybrid maize lines. Inbred maize lines need to be highly homogeneous, homozygous and reproducible to be useful as parents of commercial hybrids. There are many analytical methods available to determine the homozygotic and phenotypic stability of these inbred lines.

The oldest and most traditional method of analysis is the observation of phenotypic traits. The data is usually collected in field experiments over the life of the maize plants to be examined. Phenotypic characteristics most often observed are for traits associated with plant morphology, ear and kernel morphology, insect and disease resistance, maturity, and yield.

In addition to phenotypic observations, the genotype of a plant can also be examined. There are many laboratory-based techniques available for the analysis, comparison and characterization of plant genotype; among these are Isozyme Electrophoresis, Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), and Simple Sequence Repeats (SSRs) which are also referred to as Microsatellites.

Some of the most widely used of these laboratory techniques are Isozyme Electrophoresis and RFLPs as discussed in Lee, M., "Inbred Lines of Maize and Their Molecular Markers," *The Maize Handbook*, (Springer-Verlag, New York, Inc. 1994, at 423-432). Isozyme Electrophoresis is a useful tool in determining genetic composition, although it has relatively low number of available markers and the low number of allelic variants among maize inbreds. RFLPs have the advantage of revealing an exceptionally high degree of allelic variation in maize and the number of available markers is almost limitless. Maize RFLP linkage maps have been rapidly constructed and widely implemented in genetic studies. One such study is described in Boppenmaier, et al., "Comparisons among strains of inbreds for RFLPs", *Maize Genetics Cooperative Newsletter*, 65:1991, pg. 90. This study used 101 RFLP markers to analyze the patterns of 2 to 3 different deposits each of five different inbred lines. The inbred lines had been selfed from 9 to 12 times before being adopted into 2 to 3 different breeding programs. It was results from these 2 to 3 different breeding programs that supplied the different deposits for analysis. These five lines were maintained in the separate breeding programs by selfing or sibbing and rogueing off-type plants for an additional one to eight generations. After the RFLP analysis was completed, it was determined the five lines showed 0-2% residual heterozygosity. Although this was a relatively small study, it can be seen using RFLPs that the lines had been highly homozygous prior to the separate strain maintenance.

The production of hybrid maize lines typically comprises planting in pollinating proximity seeds of, for example, inbred maize line NP2052 and of a different inbred parent maize plant, cultivating the seeds of inbred maize line NP2052 and of said different inbred parent maize plant into plants that bear flowers, emasculating the male flowers of inbred maize line NP2052 or the male flowers of said different inbred parent maize plant to produce an emasculated maize plant, allowing cross-pollination to occur between inbred maize line NP2052 and said different inbred parent maize plant and harvesting seeds produced on said emasculated maize plant. The harvested seed are grown to produce hybrid maize plants.

Inbred maize line NP2052 can be crossed to inbred maize lines of various heterotic group (see e.g. Hallauer et al. (1988) in Corn and Corn Improvement, Sprague et al, eds, chapter 8, pages 463-564) for the production of hybrid maize lines.

TABLE I
VARIETY DESCRIPTION INFORMATION
Inbred maize line NP2052 is compared to inbred A619

	INBRED NP2052			INBRED A619		
<u>MATURITY</u>	Days	Heat		Days	Heat	
		Units			Units	
From emergence to 50% of plants in silk	66	1293.3		66	1314.3	
From emergence to 50% of plants in pollen	65	1267.8		64	1251.9	
From 10% to 90% pollen shed	003	0074.3		003	0093.5	
<u>PLANT</u>		Std Dev	Sample		Std Dev	Sample
			Size			Size
cm Plant Height (to tassel tip)	180.3	22.1	10	180.8	30.4	10
cm Ear Height (to base of top ear node)	45.1	2.7	10	47.0	13.5	10
cm Length of Top Ear Internode	9.6	2.6	10	11.6	3.7	10
Average Number of Tillers	0	0	10	0	0.0	10
Average Number of Ears per Stalk	1.2	.3	11	1.1	0.2	11
Anthocyanin of Brace Roots:	3			2		
1=Absent 2=Faint 3=Moderate 4=Dark						
<u>LEAF</u>		Std Dev	Sample		Std Dev	Sample
			Size			Size
Cm Width of Ear Node Leaf	008.0	2.6	10	008.3	2.6	10
cm Length of Ear Node Leaf	064.8	21.8	9	060.7	19.9	9
Number of leaves above top ear	6	.4	11	5	.2	11
degrees Leaf Angle	50	12.2	11	45	16.5	11
(measure from 2nd leaf above ear at						

anthesis to stalk above leaf)					
Leaf Color	02	(Munsell code 5GY 4/4)	03	(Munsell code 5GY 3/4)	
Leaf Sheath Pubescence (Rate on scale from 1=none to 9=like peach fuzz)	5		2		
Marginal Waves (Rate on scale from 1=none to 9=many)	4		5		
Longitudinal Creases (Rate on scale from 1=none to 9=many)	5		4		
<u>TASSEL</u>					
Number of Primary Lateral Branches	7	1.5	10	8	1.9
Branch Angle from Central Spike	63	40.2	11	44	10.5
Cm Tassel Length (from top leaf collar to tassel tip)	33.9	9.6	10	34.1	9.9
Pollen Shed (Rate on scale from 0=male sterile to 9=heavy shed)	7			7	
Anther Color	26	(Munsell code)		05	(Munsell code 2.5GY 8/6)
Glume Color	26	(Munsell code)		05	(Munsell code 5GY 5/8)
Bar Glumes (Glume Bands): 1=Absent 2=Present	2			2	
<u>EAR (Unhusked Data)</u>					
Silk Color (3 days after emergence)	26	(Munsell code)		05	(Munsell code 2.5GY 8/8)
Fresh Husk Color(25 days after 50% silking)		(Munsell code)			(Munsell code)
Dry Husk Color (65 days after 50 % silking)		(Munsell code)			(Munsell code)

Position of Ear at Dry Husk Stage:
 1=Upright 2=Horizontal 3=Pendent
 Husk Tightness
 (Rate on scale from 1=very loose to 9=very tight)
 Husk Extension (at harvest):
 1=Short (ears exposed) 2=Medium (<8cm)
 3=Long
 (8-10 cm beyond ear tip) 4=Very long (>10 cm)

1
 6
 3

1
 5
 2

EAR (Husked Ear Data)

Std Dev Sample
 Size

Std Dev Sample
 Size

Cm Ear Length
 mm Ear Diameter at mid-point
 gm Ear Weight
 Number of Kernel Rows
 Kernel Rows: 1=Indistinct 2=Distinct

14.0 2.1 8
 40.1 3.1 7
 81.2 17.4 6
 15 0.9 6
 2

14.8 2.0 10
 44.1 6.3 9
 90.1 27.8 9
 15 1.2 9
 2

Row Alignment:
 1=Straight 2=Slightly Curved 3=Spiral
 cm Shank Length
 Ear Taper: 1=Slight 2=Average
 3=Extreme

1
 9.0 1.1 7
 2

2
 8.7 1.9 90
 2

KERNEL (Dried)

Std Dev Sample
 Size

Std Dev Sample
 Size

mm Kernel Length
 mm Kernel Width
 mm Kernel Thickness
 % Round Kernels (Shape Grade)
 Aleurone Color Pattern:
 1=Homozygous 2=Segregating
 Aleurone Color

10.2 0.4 6
 8.2 .8 6
 5.4 .5 6
 79.9 18.9 6
 1
 18 (Munsell code

10.5 1.1 8
 8.9 1.0 8
 4.9 1.1 8
 65.4 25.6 8
 1
 18 (Munsell code)

Hard Endosperm Color	08	7.5YR 8/4) (Munsell code 7.5YR 6/8)	07	(Munsell code 7.5YR 7/10)
Endosperm Type: 1=Sweet (su1) 2=Extra Sweet (sh2) 3=Normal Starch	3		3	
Gm Weight per 100 Kernels (unsized sample)	31.0		31.1	
<u>COB</u>		Std Dev Sample Size		Std Dev Sample Size
mm Cob Diameter at mid-point	27.1	1.2 8	27.9	2.3 9
Cob Color	19	(Munsell code)	19	(Munsell code)
<u>DISEASE RESISTANCE</u> (1=most susceptible to 9=most resistant)				
Eye Spot (Kabatiella zeae)	6		4	
Northern Leaf Blight	7	Mixed Inoc.	3	Mixed Inoc.
Gray Leaf Spot	5		5	
Common Rust	7		6	
<u>INSECT RESISTANCE</u> (Rate from 1=most susceptible to 9=most resistant)				
European Corn Borer(Osstrinia nubilalis) 1 st Generation (Typically Whorl Leaf Feeding)	6		7	
2 nd Generation Corn Borer	2		4	
<u>AGRONOMIC TRAITS</u>				
Stay Green (at 65 days after anthesis) (rate on scale from 1=worst to 9=excellent)	6		5	
% Dropped Ears (at 65 days after anthesis)	0		0	
% Pre-anthesis Brittle snapping	0		0	

% Pre-anthesis Root Lodging	0	0
% Post-anthesis Root Lodging (at 65 days after anthesis)	0	0
Kg/ha Yield of Inbred Per Se (at 12-13% grain moisture)	5067.0	3398.9

In interpreting the foregoing color designations, reference may be made to the Munsell Glossy Book of Color, a standard color reference. Color codes: 1. light green, 2. medium green, 3. dark green, 4. very dark green 5. green-yellow, 6. pale yellow, 7. yellow, 8. yellow-orange 9. salmon, 10. pink-orange, 11. pink 12. light red, 13. cherry red. 14. red, 15. red and white, 16. pale purple, 17. purple, 18. colorless, 19. white, 20. white capped, 21. buff, 22. tan, 23. brown, 24. bronze, 25. variegated, 26. other.

There are clear differences between the above two inbreds. For example, the inbred line NP2052 silk is green-yellow (Munsell code 2.5GY 8/8) with pale purple shaded ends 5RP 4/10) and A619 is green-yellow (Munsell code 2.5GY 8/8).

Some of the more pronounced differences between NP2052 and A619 occur in the tassel. The NP2052 tassel appearance is not as compact with tassel branches having a larger angle from the central spike giving it a "droopy" look as compared to A619. The A619 tassel is compact with upright shorter tassel branches. The NP2052 anther color is pale yellow (Munsell code 5Y 8/8) with pale purple shading (Munsell code 5RP 4/10). The anther color of A619 is green-yellow (Munsell code 2.5GY 8/6). NP2052 has a glume color of medium green (Munsell code 5GY 4/6) with pale purple coloration (Munsell code 5RP 4/10). The glume of A619 is green-yellow (Munsell code 5GY 5/8). The glume band of NP2052 has a pale purple shade. The glume band of A619 is light green.

The leaf color of A619 is darker than NP2052. The leaf color for A619 is dark green (Munsell code 5GY 3/4). The leaf color for NP2052 is medium green (Munsell code 5GY 4/4).

There are several differences in disease and insect ratings when comparing NP2052 and A619. NP2052 has a rating of 7 for Northern Leaf Blight – Mixed Inoculum and A619 is rated a 3. The Eyespot rating for NP2052 is a 6 as compared to a rating of 4 for A619. The Common Rust rating for NP2052 is 7 and A619 is a 6. The First Brood Corn Borer rating for NP2052 is a 6 with A619 rated as a 7. The Second Brood Corn Borer rating for NP2052 is a 2 as compared to a 4 for A619.

The yield of NP2052 is greater than A619. The Kg/ha yield of NP2052 is 5067.0. The yield of A619 is 3398.9 Kg/ha.

Inbred maize line NP2052 is a yellow dent maize inbred that can be used as a male or female in crosses for producing first generation F1 maize hybrids. Inbred maize line NP2052 is best adapted

to the Northern Cornbelt regions of Canada and the United States. Hybrids made with NP2052 consistently yielded above trial mean in 1998 and 1999. The inbred can be used to produce hybrids from approximately 85-105 relative maturity based on the Comparative Relative Maturity Rating System for harvest moisture of grain. Inbred maize line NP2052 demonstrates reliable late season plant health and average pollen shed, with 7-8 lateral branches. In hybrid combinations, NP2052 demonstrates high yield for its maturity, sound seedling vigor, average plant stature, improved stalk strength, rapid drydown, and has an attractive plant appearance. In hybrid combination, NP2052 also demonstrates acceptable resistance to various stalk rots, good resistance to Northern Leaf Blight, good resistance to Eyespot, good resistance to Common Rust, and good resistance to First Brood Corn Borer. For its area of adaptation, NP2052 demonstrates high yields, resistance to stalk diseases, acceptable late season intactness, and reliable seedling vigor.

Origin and Breeding History of Corn Inbred Line NP2052

Inbred corn line NP2052 was derived from self pollinating the hybrid Pioneer Brand 3751. After development of the S_0 (or F_2) population, the breeding method was simple pedigree ear-to-row.

The details of the development of inbred line NP2052 are as follows:

1988 Stanton, MN Pioneer Brand 3751 was self-pollinated to produce S_0 seed.

1988 Kauai, Hawaii S_0 plants were self pollinated to produce the S_1 generation. Two hundred fourteen S_1 ears were selected from individual S_0 plants based upon plant quality, root strength, ear size, and resistance to diseases.

1989 Stanton, MN Ears of the two hundred fourteen selected S_1 families were grown, observed, and self-pollinated to produce the S_2 generation. Phenotypic selection of these S_1 families was based upon plant quality, synchrony of pollen shed and silk emergence, root strength, ear size, and resistance to disease. Testcross pollinations of the S_1 families were also made.

1990 Stanton, MN Ear rows of the selected S_2 families were grown and observed. The S_2 families were also self-pollinated to produce the S_3 generation. Selection of the S_2 families was based upon the performance of the S_1 testcrosses for grain yield, maturity, and general quality. These testcrosses were grown at several locations. Phenotypic selection of the S_2 families was based upon plant quality, synchrony of pollen shed and silk emergence, root strength, ear size, and resistance to disease. Testcrosses of S_2 families were also made.

1990 Puerto Rico Ear rows of selected S_3 families were grown, observed, and self-pollinated to produce the S_4 generation. Selection of S_3 families was based upon performance of the S_2 testcrosses for grain yield, maturity and general quality. These testcrosses were grown at several locations. Phenotypic selection of S_3 families was based upon plant quality, synchrony of pollen shed

and silk emergence, root strength, ear size, and resistance to disease. Testcrosses of the S_3 families were also made.

1991 Stanton, MN Ear rows of selected S_4 families were grown, observed, and self-pollinated to produce the S_5 generation. Selection of the S_4 families was based upon performance of the S_2 and S_3 testcrosses for grain yield, maturity and general quality. These testcrosses were grown at several locations. Phenotypic selection of the S_4 families was based upon synchrony of pollen shed and silk emergence, root strength, ear size, and resistance to disease. Testcrosses of the S_4 families were also made.

1991 Kauai, Hawaii Ear rows of selected S_5 families were grown, observed, and self-pollinated to produce the S_6 generation.

1992 Stanton, MN Ear rows of each selected S_6 family were grown, observed and self-pollinated to produce individual S_7 ears. Selection of the S_6 families was based upon the performance of the S_4 testcrosses for grain yield, maturity, and general quality. These testcrosses were grown at several locations. Phenotypic selection of the S_6 families was based upon synchrony of pollen shed and silk emergence, root strength, ear size, and resistance to disease.

1993 Stanton, MN Rows of each selected S_7 individual ear culture were grown, observed, and self-pollinated to produce "breeder's seed." These families were closely evaluated and selected for uniformity of anther and silk color, plant and ear height, and other characteristics. S_8 individual ears were saved from one S_7 individual ear culture.

1993 Kauai, Hawaii Rows of each selected S_8 individual ear were grown, observed and self-pollinated to produce additional "breeder's seed." Plants were closely evaluated for uniformity of anther and silk color, plant and ear height, and other characteristics. Isozyme testing confirmed the purity of this inbred line. S_9 individual ears were saved from one S_8 individual ear family.

1994 Stanton, MN Rows of the selected S_9 individual ears were grown, observed, and self-pollinated to produce additional "breeder's seed." Plants were closely evaluated for uniformity of anther and silk color, plant and ear height, and other characteristics. Isozyme testing confirmed the purity of this inbred line.

From 1994 to the present this inbred line has been observed in Stanton, MN and other locations. No phenotypic or isozymic variants have been observed. The variety NP2052 has been uniform and stable.

The invention also encompasses plants of inbred maize line NP2052 and parts thereof further comprising one or more specific, single gene traits which have been introgressed into inbred maize line NP2052 from another maize line. Preferably, one or more new traits are transferred to inbred maize line NP2052, or, alternatively, one or more traits of inbred maize line NP2052 are altered or

substituted. The transfer (or introgression) of the trait(s) into inbred maize line NP2052 is for example achieved by recurrent selection breeding, for example by backcrossing. In this case, inbred maize line NP2052 (the recurrent parent) is first crossed to a donor inbred (the non-recurrent parent) that carries the appropriate gene(s) for the trait(s) in question. The progeny of this cross is then mated back to the recurrent parent followed by selection in the resultant progeny for the desired trait(s) to be transferred from the non-recurrent parent. After three, preferably four, more preferably five or more generations of backcrosses with the recurrent parent with selection for the desired trait(s), the progeny will be heterozygous for loci controlling the trait(s) being transferred, but will be like the recurrent parent for most or almost all other genes (see, for example, Poehlman & Sleper (1995) *Breeding Field Crops*, 4th Ed., 172-175; Fehr (1987) *Principles of Cultivar Development*, Vol. 1: Theory and Technique, 360-376).

sub 21. The laboratory-based techniques described above, in particular RFLP and SSR, are routinely used in such backcrosses to identify the progenies having the highest degree of genetic identity with the recurrent parent. This permits to accelerate the production of inbred maize lines having at least 90%, preferably at least 95%, more preferably at least 99% genetic identity with the recurrent parent, yet more preferably genetically identical to the recurrent parent, and further comprising the trait(s) introgressed from the donor parent. Such determination of genetic identity is based on molecular markers used in the laboratory-based techniques described above. Such molecular markers are for example those known in the art and described in Boppenmaier, et al., "Comparisons among strains of inbreds for RFLPs", *Maize Genetics Cooperative Newsletter* (1991) 65, pg. 90, or those available from the University of Missouri database and the Brookhaven laboratory database (see <http://www.agron.missouri.edu>). The last backcross generation is then selfed to give pure breeding progeny for the gene(s) being transferred. The resulting plants have essentially all of the morphological and physiological characteristics of inbred maize line NP2052, in addition to the single gene trait(s) transferred to the inbred. Preferably, the resulting plants have all of the morphological and physiological characteristics of inbred maize line NP2052, in addition to the single gene trait(s) transferred to the inbred. The exact backcrossing protocol will depend on the trait being altered to determine an appropriate testing protocol. Although backcrossing methods are simplified when the trait being transferred is a dominant allele, a recessive allele may also be transferred. In this instance it may be necessary to introduce a test of the progeny to determine if the desired trait has been successfully transferred.

Many traits have been identified that are not regularly selected for in the development of a new inbred but that can be improved by backcrossing techniques or genetic transformation. Examples of traits transferred to inbred maize line NP2052 include, but are not limited to, waxy starch,

herbicide tolerance, resistance for bacterial, fungal, or viral disease, insect resistance, enhanced nutritional quality, improved performance in an industrial process, altered reproductive capability, such as male sterility or male fertility, yield stability and yield enhancement. Other traits transferred to inbred maize line NP2052 are for the production of commercially valuable enzymes or metabolites in plants of inbred maize line NP2052.

Traits transferred to maize inbred line NP2052 are naturally occurring maize traits, which are preferably introgressed into inbred maize line NP2052 by breeding methods such as backcrossing, or are heterologous transgenes, which are preferably first introduced into a maize line by genetic transformation using genetic engineering and transformation techniques well known in the art, and then introgressed into inbred line NP2052. Alternatively a heterologous trait is directly introduced into inbred maize line NP2052 by genetic transformation. Heterologous, as used herein, means of different natural origin or represents a non-natural state. For example, if a host cell is transformed with a nucleotide sequence derived from another organism, particularly from another species, that nucleotide sequence is heterologous with respect to that host cell and also with respect to descendants of the host cell which carry that gene. Similarly, heterologous refers to a nucleotide sequence derived from and inserted into the same natural, original cell type, but which is present in a non-natural state, e.g. a different copy number, or under the control of different regulatory sequences. A transforming nucleotide sequence may comprise a heterologous coding sequence, or heterologous regulatory sequences. Alternatively, the transforming nucleotide sequence may be completely heterologous or may comprise any possible combination of heterologous and endogenous nucleic acid sequences.

A transgene introgressed into maize inbred line NP2052 typically comprises a nucleotide sequence whose expression is responsible or contributes to the trait under the control of a promoter appropriate for the expression of the nucleotide sequence at the desired time in the desired tissue or part of the plant. Constitutive or inducible promoters are used. The transgene may also comprise other regulatory elements such as for example translation enhancers or termination signals. In a preferred embodiment, the nucleotide sequence is the coding sequence of a gene and is transcribed and translated into a protein. In another preferred embodiment, the nucleotide sequence encodes an antisense RNA, a sense RNA that is not translated or only partially translated, a t-RNA, a r-RNA or a sn-RNA.

Where more than one trait are introgressed into inbred maize line NP2052, it is preferred that the specific genes are all located at the same genomic locus in the donor, non-recurrent parent, preferably, in the case of transgenes, as part of a single DNA construct integrated into the donor's genome. Alternatively, if the genes are located at different genomic loci in the donor, non-recurrent

parent, backcrossing allows to recover all of the morphological and physiological characteristics of inbred maize line NP2052 in addition to the multiple genes in the resulting maize inbred line.

The genes responsible for a specific, single gene trait are generally inherited through the nucleus. Known exceptions are, e.g. the genes for male sterility, some of which are inherited cytoplasmically, but still act as single gene traits. In a preferred embodiment, a heterologous transgene to be transferred to maize inbred line NP2052 is integrated into the nuclear genome of the donor, non-recurrent parent. In another preferred embodiment, a heterologous transgene to be transferred to maize inbred line NP2052 is integrated into the plastid genome of the donor, non-recurrent parent. In a preferred embodiment, a plastid transgene comprises one gene transcribed from a single promoter or two or more genes transcribed from a single promoter.

In a preferred embodiment, a transgene whose expression results or contributes to a desired trait to be transferred to maize inbred line NP2052 comprises a virus resistance trait such as, for example, a MDMV strain B coat protein gene whose expression confers resistance to mixed infections of maize dwarf mosaic virus and maize chlorotic mottle virus in transgenic maize plants (Murry et al. *Biotechnology* (1993) 11:1559-64). In another preferred embodiment, a transgene comprises a gene encoding an insecticidal protein, such as, for example, a crystal protein of *Bacillus thuringiensis* or a vegetative insecticidal protein from *Bacillus cereus*, such as VIP3 (see for example Estruch et al. *Nat Biotechnol* (1997) 15:137-41). In a preferred embodiment, an insecticidal gene introduced into maize inbred line NP2052 is a Cry1Ab gene or a portion thereof, for example introgressed into maize inbred line NP2052 from a maize line comprising a Bt-11 event as described in U.S. Patent No. 6,114,608, which is incorporated herein by reference, or from a maize line comprising a 176 event as described in Koziel *et al.* (1993) *Biotechnology* 11: 194-200. In yet another preferred embodiment, a transgene introgressed into maize inbred line NP2052 comprises a herbicide tolerance gene. For example, expression of an altered acetohydroxyacid synthase (AHAS) enzyme confers upon plants tolerance to various imidazolinone or sulfonamide herbicides (U.S. Patent No. 4,761,373, incorporated herein by reference). In another preferred embodiment, a non-transgenic trait conferring tolerance to imidazolinones is introgressed into maize inbred line NP2052 (e.g. a "IT" or "IR" trait). U.S. Patent No. 4,975,374, incorporated herein by reference, relates to plant cells and plants containing a gene encoding a mutant glutamine synthetase (GS) resistant to inhibition by herbicides that are known to inhibit GS, e.g. phosphinothricin and methionine sulfoximine. Also, expression of a *Streptomyces bar* gene encoding a phosphinothricin acetyl transferase in maize plants results in tolerance to the herbicide phosphinothricin or glufosinate (U.S. Patent No. 5,489,520, incorporated herein by reference). U.S. Patent No. 5,013,659, incorporated herein by reference, is directed to plants that express a mutant acetolactate synthase (ALS) that renders the plants resistant to

inhibition by sulfonylurea herbicides. U.S. Patent No. 5,162,602, incorporated herein by reference, discloses plants tolerant to inhibition by cyclohexanedione and aryloxyphenoxypropanoic acid herbicides. The tolerance is conferred by an altered acetyl coenzyme A carboxylase (ACCase). U.S. Patent No. 5,554,798, incorporated herein by reference, discloses transgenic glyphosate tolerant maize plants, which tolerance is conferred by an altered 5-enolpyruvyl-3-phosphoshikimate (EPSP) synthase gene. U.S. Patent No. 5,804,425, incorporated herein by reference, discloses transgenic glyphosate tolerant maize plants, which tolerance is conferred by an EPSP synthase gene derived from *Agrobacterium tumefaciens* CP-4 strain. Also, tolerance to a protoporphyrinogen oxidase inhibitor is achieved by expression of a tolerant protoporphyrinogen oxidase enzyme in plants (U.S. Patent No. 5,767,373, which is incorporated herein by reference). Another trait transferred to inbred maize line NP2052 confers tolerance to an inhibitor of the enzyme hydroxyphenylpyruvate dioxygenase (HPPD) and transgenes conferring such trait are, for example, described in WO 9638567, WO 9802562, WO 9923886, WO 9925842, WO 9749816, WO 9804685 and WO 9904021.

In a preferred embodiment, a transgene transferred to maize inbred line NP2052 comprises a gene conferring tolerance to a herbicide and at least another nucleotide sequence encoding another trait, such as for example, an insecticidal protein. Such combination of single gene traits is for example a Cry1Ab gene and a *bar* gene.

Specific transgenic events introgressed into maize inbred line NP2052 are found at http://www.aphis.usda.gov/bbep/bp/not_reg.html. For example, introgressed from glyphosate tolerant event GA21 (9709901p), glyphosate tolerant/Lepidopteran insect resistant event MON 802 (9631701p), Lepidopteran insect resistant event DBT418 (9629101p), male sterile event MS3 (9522801p), Lepidopteran insect resistant event Bt11 (9519501p), phosphinothricin tolerant event B16 (9514501p), Lepidopteran insect resistant event MON 80100 (9509301p), phosphinothricin tolerant events T14, T25 (9435701p), Lepidopteran insect resistant event 176 (9431901p).

The introgression of a Bt11 event into a maize line, such as maize inbred line NP2052, by backcrossing is exemplified in U.S. Patent No. 6,114,608, and the present invention is directed to methods of introgressing a Bt11 event into maize inbred line NP2052 using for example the markers described in U.S. Pat. No. 6,114,608 and to resulting maize lines.

Direct selection may be applied where the trait acts as a dominant trait. An example of a dominant trait is herbicide tolerance. For this selection process, the progeny of the initial cross are sprayed with the herbicide prior to the backcrossing. The spraying eliminates any plant which do not have the desired herbicide tolerance characteristic, and only those plants that have the herbicide tolerance gene are used in the subsequent backcross. This process is then repeated for the additional backcross generations.

This invention also is directed to methods for producing a maize plant by crossing a first parent maize plant with a second parent maize plant wherein either the first or second parent maize plant is a maize plant of inbred line NP2052 or a maize plant of inbred line NP2052 further comprising one or more single gene traits. Further, both first and second parent maize plants can come from the inbred maize line NP2052 or an inbred maize plant of NP2052 further comprising one or more single gene traits. Thus, any such methods using the inbred maize line NP2052 or an inbred maize plant of NP2052 further comprising one or more single gene traits are part of this invention: selfing, backcrosses, hybrid production, crosses to populations, and the like. All plants produced using inbred maize line NP2052 or inbred maize plants of NP2052 further comprising one or more single gene traits as a parent are within the scope of this invention. Advantageously, inbred maize line NP2052 or inbred maize plants of NP2052 further comprising one or more single gene traits are used in crosses with other, different, maize inbreds to produce first generation (F1) maize hybrid seeds and plants with superior characteristics.

In a preferred embodiment, seeds of inbred maize line NP2052 or seeds of inbred maize plants of NP2052 further comprising one or more single gene traits are provided as an essentially homogeneous population of inbred corn seeds. Essentially homogeneous populations of inbred seed are those that consist essentially of the particular inbred seed, and are generally purified free from substantial numbers of other seed, so that the inbred seed forms between about 90% and about 100% of the total seed, and preferably, between about 95% and about 100% of the total seed. Most preferably, an essentially homogeneous population of inbred corn seed will contain between about 98.5%, 99%, 99.5% and about 100% of inbred seed, as measured by seed grow outs. The population of inbred corn seeds of the invention is further particularly defined as being essentially free from hybrid seed. The inbred seed population may be separately grown to provide an essentially homogeneous population of plants of inbred maize line NP2052 or inbred maize plants of NP2052 further comprising one or more single gene traits.

As used herein, the term "plant" includes plant cells, plant protoplasts, plant cell tissue cultures from which maize plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants, such as embryos, pollen, ovules, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk, seeds and the like.

Duncan, Williams, Zehr, and Widholm, *Planta* (1985) 165:322-332 reflects that 97% of the plants cultured that produced callus were capable of plant regeneration. Subsequent experiments with both inbreds and hybrids produced 91% regenerable callus that produced plants. In a further study in 1988, Songstad, Duncan & Widholm in *Plant Cell Reports* (1988), 7:262-265 reports several media additions that enhance regenerability of callus of two inbred lines. Other published reports also

indicated that "nontraditional" tissues are capable of producing somatic embryogenesis and plant regeneration. K. P. Rao, et al., Maize Genetics Cooperation Newsletter, 60:64-65 (1986), refers to somatic embryogenesis from glume callus cultures and B. V. Conger, et al., Plant Cell Reports, 6:345-347 (1987) indicates somatic embryogenesis from the tissue cultures of maize leaf segments. Thus, it is clear from the literature that the state of the art is such that these methods of obtaining plants are, and were, "conventional" in the sense that they are routinely used and have a very high rate of success.

Tissue culture procedures of maize are described in Green and Rhodes, "Plant Regeneration in Tissue Culture of Maize," Maize for Biological Research (Plant Molecular Biology Association, Charlottesville, Va. 1982, at 367-372) and in Duncan, et al., "The Production of Callus Capable of Plant Regeneration from Immature Embryos of Numerous Zea mays Genotypes," 165 Planta 322-332 (1985). Thus, another aspect of this invention is to provide cells that upon growth and differentiation produce maize plants having the physiological and morphological characteristics of inbred maize line NP2052. In a preferred embodiment, cells of inbred maize line NP2052 are transformed genetically, for example with one or more genes described above, for example by using a transformation method described in U.S. Pat. No. 6,114,608, and transgenic plants of inbred maize line NP2052 are obtained and used for the production of hybrid maize plants.

Maize is used as human food, livestock feed, and as raw material in industry. The food uses of maize, in addition to human consumption of maize kernels, include both products of dry- and wet-milling industries. The principal products of maize dry milling are grits, meal and flour. The maize wet-milling industry can provide maize starch, maize syrups, and dextrose for food use. Maize oil is recovered from maize germ, which is a by-product of both dry- and wet-milling industries.

Maize, including both grain and non-grain portions of the plant, is also used extensively as livestock feed, primarily for beef cattle, dairy cattle, hogs, and poultry. Industrial uses of maize include production of ethanol, maize starch in the wet-milling industry and maize flour in the dry-milling industry. The industrial applications of maize starch and flour are based on functional properties, such as viscosity, film formation, adhesive properties, and ability to suspend particles. The maize starch and flour have application in the paper and textile industries. Other industrial uses include applications in adhesives, building materials, foundry binders, laundry starches, explosives, oil-well muds, and other mining applications. Plant parts other than the grain of maize are also used in industry: for example, stalks and husks are made into paper and wallboard and cobs are used for fuel and to make charcoal.

The seed of inbred maize line NP2052 or of inbred maize line NP2052 further comprising one or more single gene traits, the plant produced from the inbred seed, the hybrid maize plant

produced from the crossing of the inbred, hybrid seed, and various parts of the hybrid maize plant can be utilized for human food, livestock feed, and as a raw material in industry.

The present invention therefore also discloses an agricultural product comprising a plant of the present invention or derived from a plant of the present invention. The present invention also discloses an industrial product comprising a plant of the present invention or derived from a plant of the present invention. The present invention further discloses methods of producing an agricultural or industrial product comprising planting seeds of the present invention, growing plant from such seeds, harvesting the plants and processing them to obtain an agricultural or industrial product.

PERFORMANCE EXAMPLES OF MAIZE INBRED LINE NP2052

In the examples that follow, the traits and characteristics of inbred maize line NP2052 are presented as an comparison between it and inbred line A619. The data presented is for key characteristics and traits and is reported from experiments where the compared lines were grown side-by-side.

The data in Table 2 show, for example, that the pollen shed of variety NP2052 is significantly shorter than A619. The length of shed for NP2052 is 74 heat units as compared to 94 for A619. Table 2 also shows that the length of the top ear internode is significantly shorter for NP2052 compared to A619, and that the length of the ear node leaf of NP2052 is significantly longer than A619.

TABLE 2

	H.U. 10% to 90% Pollen Shed	Length of Top Ear Internode (cm)	Length of Ear Node Leaf (cm)	Leaf Sheath Pubescence (1-9 rating)
Grand Mean	84	11	63	3
Trails w/data	5	6	5	6
LSD (= 95% confidence level)	16	1	3	2
CV%	27	12	4	35
Probability %	2	0	0	1
A619	94	12	61	2
NP2052	74	10	65	5

TABLE 3
VARIETY COMPARISON

	STDP	LRTL	ERTLN	ERTLP	GRSN	GRSNP	DROPN
Grand Me	100	20	0	0	0	0	0
Trials w/da	6	3	2	2	2	2	2
LSD	0	161	0	0	0	0	
CV%	0	228					
Probability		42					
A619	100	39	0	0	0	0	0
NP2052	100	1	0	0	0	0	0
	PLHCN	ERHCN	HU1SN	HU5SN	HU9SN	HUPSN	STGRP
Grand Me	171	44	1259	1314	1313	1283	6
Trials w/da	3	3	3	3	3	3	3
LSD	46	15	9	354	494	12	0
CV%	6	29	1	2	2	1	0
Probability	67	49	0	73	51	0	
A619	168	41	1227	1308	1296	1257	6
NP2052	174	46	1305	1322	1338	1322	7
	HUPLN	PLQUR	EARPN	SHLNN	HSKCR	OTHIN	
Grand Me	35	6	1	8	3	0	
Trials w/da	3	3	3	3	2	3	
LSD	22	1	0	3	6	0	
CV%	65	22	22	33	0		
Probability	0	9	17	99	20		
A619	55	7	1	8	3	0	
NP2052	15	6	1	8	4	0	
	EINDN	LTEIN	ERLWN	ERLLN	LAERN	ANGBN	LTBRN
Grand Me	5	11	9	71	4	41	8
Trials w/da	4	3	3	3	3	3	3
LSD		6	2	3	7	8	141
CV%		12	20	3	14	18	21
Probability		31	77	0	71	16	49
A619	2	12	10	69	5	38	41
NP2052	3	10	9	73	4	44	69
	TBANN	LTASN	ERLNN	ERDIN	EWGTN	KRRWN	KRLNN
Grand Me	55	39	12	37	79	13	11
Trials w/da	3	3	3	3	3	3	3
LSD	141	2	1	4	14	2	1
CV%	21	5	8	9	16	10	5
Probability	49	77	6	2	0	69	6
A619	41	39	13	39	89	13	11
NP2052	69	39	11	34	68	13	10

	KRWDN	KRDPN	KRPPN	K100N	COBDN	APBRR	LSPUR
Grand Me	8	5	69	31	26	3	3
Trials w/da	3	3	3	3	3	3	3
LSD	3	3	92	4	2	1	6
CV%	8	13	18	10	8	21	43
Probability	37	76	50	42	10	18	14
A619	9	5	61	32	27	2	2
NP2052	8	5	78	30	25	3	5

	MLWVR	LFLCR	HSKTR	HEPSR	KRRWR	KRNAR	ETAPR	TILLN
Grand Me	5	5	7	1	2	2	2	0.0095
Trials w/da	3	3	2	3	3	3	3	3
LSD	5	1	6	0	0	0	0	0.7874
CV%	35	34	0	31	0	14	7	7230
Probability	23	24	20	100	0	97	0	96.19
A619	6	5	6	1	2	2	2	0.019
NP2052	4	5	8	1	2	2	2	0

DEPOSIT

546 2³ Applicants have made a deposit of at least 2500 seeds of Inbred Maize Line NP2052 with the American Type Culture Collection (ATCC), Manassas, Virginia, 20110-2209 U.S.A., ATCC Deposit No: _____. This deposit of the Inbred Maize Line NP2052 will be maintained in the ATCC depository, which is a public depository, for a period of 30 years, or 5 years after the most recent request, or for the effective life of the patent, whichever is longer, and will be replaced if it becomes nonviable during that period. Additionally, Applicants have satisfied all the requirements of 37 C.F.R. §§1.801-1.809, including providing an indication of the viability of the sample. Applicants impose no restrictions on the availability of the deposited material from the ATCC; however, Applicants have no authority to waive any restrictions imposed by law on the transfer of biological material or its transportation in commerce. Applicants do not waive any infringement of its rights granted under this patent or under the Plant Variety Protection Act (7 USC 2321 et seq.).

The foregoing invention has been described in detail by way of illustration and example for purposes of clarity and understanding. However, it will be obvious that certain changes and modifications such as single gene modifications and mutations, somaclonal variants, variant individuals selected from large populations of the plants of the instant inbred and the like may be practiced within the scope of the invention, as limited only by the scope of the appended claims.